AD	1		

Award Number: DAMD17-02-1-0470

TITLE: Intervention Study of Flaxseed in Postmenopausal Women: Effects on

Hormonal Biomarkers of Breast Cancer Risk

PRINCIPAL INVESTIGATOR: Susan R. Sturgeon, Ph.D.

CONTRACTING ORGANIZATION: University of Massachusetts

Amherst, MA 01003-9272

REPORT DATE: January 2006

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;

Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188		
Public reporting burden for this collection of information is data needed, and completing and reviewing this collection this burden to Department of Defense, Washington Headq 4302. Respondents should be aware that notwithstanding	estimated to average 1 hour per responsion of information. Send comments regularters Services, Directorate for Infoliany other provision of law, no perso	oonse, including the time for revie arding this burden estimate or ar rmation Operations and Reports n shall be subject to any penalty	ny other aspect of this (0704-0188), 1215 J	OIVID TVO. 07/04-01/08 arching existing data sources, gathering and maintaining the collection of information, including suggestions for reducing efferson Davis Highway, Suite 1204, Arlington, VA 22202-with a collection of information if it does not display a currently		
valid OMB control number. PLEASE DO NOT RETURN Y 1. REPORT DATE (DD-MM-YYYY)	OUR FORM TO THE ABOVE ADDI 2. REPORT TYPE	RESS.	3	. DATES COVERED (From - To)		
01-01-2006	Final			JUL 2002 - 30 DEC 2005		
4. TITLE AND SUBTITLE Intervention Study of Flaxseed in P	ostmenopausal Wome	en: Effects on	5.	a. CONTRACT NUMBER		
Hormonal Biomarkers of Breast Ca	ncer Risk		5	b. GRANT NUMBER		
				AMD17-02-1-0470		
			5	c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S) Susan R. Sturgeon, Ph.D.			5	d. PROJECT NUMBER		
			5	e. TASK NUMBER		
Consiliractives and Quahaalah waas			5	f. WORK UNIT NUMBER		
E-mail: ssturgeon@schoolph.umas 7. PERFORMING ORGANIZATION NAME	(S) AND ADDRESS(ES)		R	PERFORMING ORGANIZATION REPORT		
University of Massachusetts Amherst, MA 01003-9272	(3) AND ADDICESS(ES)			NUMBER		
9. SPONSORING / MONITORING AGENCY U.S. Army Medical Research and M Fort Detrick, Maryland 21702-5012	Materiel Command	S(ES)		D. SPONSOR/MONITOR'S ACRONYM(S)		
			1	1. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION / AVAILABILITY STAT Approved for Public Release; Distri						
13. SUPPLEMENTARY NOTES						
14. ABSTRACT The objective of this study is to determ a beneficial effect on important of usual diet plus 7.5 grams per day of raw flaxseed. Serum and urinary baseline, 7 weeks and 13 weeks. Testradiol after six and 12 weeks of significant. Among overweight/obes substantial declines in serum levels flaxseed diet. Serum levels of sex hoverweight/obese women. Addition metabolites associated with flaxseed	t hormonal biomarkers y of raw flaxseed, follow levels of hormonal biomarkers here were small decling flaxseed ingestion among se women (body mass s of testosterone, estronormone binding globulat analyses are under	s of breast cancer rispowed by 6 weeks of comarkers of breast ones in serum levels ong all women (n=4 s index > 25kg/m ₂) (one and estradiol as all in were essentially	sk. Participar usual diet plucancer risk watestosterone (17) but these (17), there is sociated with a unchanged	nts received 6 weeks us 15 grams per day were assessed at , estrone and were not statistically were more n consumption of a in normal and		
15. SUBJECT TERMS No subject terms provided.						
16. SECURITY CLASSIFICATION OF:		17. I IMITATION	18 NUMBER	19a, NAME OF RESPONSIBLE PERSON		

OF ABSTRACT

UU

a. REPORT

b. ABSTRACT

U

c. THIS PAGE

U

OF PAGES

8

USAMRMC

19b. TELEPHONE NUMBER (include area code)

Table of Contents

Cover	1
SF 298	2
Introduction	4
Body	4
Key Research Accomplishments	5
Conclusions	6
Reportable Outcomes	6
References	6
Appendices	7

Introduction: It has been hypothesized that plant-derived estrogens (phytoestrogens) play an important role in the etiology of breast cancer. One class of phytoestrogens known as lignans are present in a variety of fruits, vegetables, and grains, but flaxseed is by far, the richest source. Laboratory and cross-sectional studies indicate that lignans are favorably associated with important hormonal biomarkers of breast cancer risk, including elevated serum levels of sex-hormone binding globulin and lower serum levels of estradiol and testosterone. The objective of this study is to determine if flaxseed supplementation to usual diet in postmenopausal women has a beneficial effect on important hormonal biomarkers of breast cancer risk. The specific aims of this study are to determine if (1) flaxseed intake lowers serum levels of estradiol and testosterone and increases serum levels of sex-hormone binding globulin (SHBG); (2) flaxseed intake increases the urinary excretion ratio of 2/4-OHE1 (OHE1:hydroxyestrone) and 2/16α-OHE1, and decreases the urinary excretion ratio of genotoxic estrogen metabolites to total estrogens ((16α-OHE1 + 4-OHE2 (OHE2:hydroxestradiol) + 4OHE1))/total estrogens; and (3) higher doses and longer duration of flaxseed intake results in greater changes to the aforementioned variables. Participants in this study will be volunteers recruited from the University of Massachusetts area. The pre-post intervention study will involve a total of 50 postmenopausal women. Participants will receive 6 weeks of usual diet plus 7.5 grams per day of raw flaxseed, followed by 6 weeks of usual diet plus 15 grams per day of raw flaxseed. Serum and urinary levels of hormonal biomarkers of breast cancer risk will be assessed at baseline, 7 weeks, and 13 weeks. The project will establish whether flaxseed may be a useful dietary strategy in the prevention of breast cancer. After discussion with the CORE, no further justification regarding the dosing schedule is necessary because this information is detailed in the protocol.

Body:

Tasks 1-4: Study participant recruitment and collection of biological specimens is complete. A total of 47 eligible study subjects successfully completed the study.

Task 5: Laboratory analyses of serum hormone levels were completed by the University of Southern California Reproductive Endocrine Laboratory. Duplicate quality control specimens from two postmenopausal women were inserted into each of the four batches. Between batch coefficients of variations ranged from 8.1% to 9.4% for testosterone, 11.5% to 12.9% for estrone, 14.3% to 15.5% for estardiol; and 3.9% to 6.4% for SHBG, indicating that laboratory assays were of acceptable quality. Urine specimens were sent to Dr. Mindy Kurzer (University of Minnesota), and laboratory analyses of estrogen metabolites and lignans have been completed.

Task 6-7: Table 1 describes the preliminary findings of the association between dietary flaxseed and changes in serum hormone levels. Serum levels of estrone, estradiol and tesosterone appeared to decline after flaxseed supplemention although these changes were not statistically significant and there was no evidence that levels declined more at increasing levels of dose and duration of intake. Serum levels of SHBG were unchanged. Adjustment for changes in other factors, including weight, physical activity and dietary factors had little impact on these findings.

Table 1: Change in hormone outcomes between each intervention visit

	Change from Baseline	Change from Baseline
	to	to
Hormone	Follow-up 1 ^{ac}	Follow-up 2 ^{bd}
Testosterone (pg/ml)	-9.91 ± 6.92	-4.65 ± 6.54
95% CI	(-23.83, 4.01)	(-17.81, 8.52)
p-value	0.16	0.48
Estrone (pg/ml)	-2.60 ± 2.06	-3.26 ± 2.23
95% CI	(-6.74, 1.54)	(-7.75, 1.22)
p-value	0.21	0.15
Estradiol (pg/ml)	-2.71 ± 1.93	-4.36 ± 4.10
95% CI	(-6.59, 1.17)	(-12.80, 3.89)
p-value	0.17	0.29
SHBG (nmol/ml)	0.59 ± 1.54	0.04 ± 1.79
95% CI	(-2.51, 3.67)	(-3.57, 3.64)
p-value	0.71	0.98

^an = 48; ^bn = 47; ^c(Follow-up 1 – Baseline); ^d(Follow-up 2 – Baseline); ^e(Follow-up 2 – Follow-up 1)

Among overweight/obese women (body mass index $\geq 25 \text{kg/m}^2$) (n=23), there were more substantial declines in serum levels of testosterone, estrone and estradiol associated with consumption of a flaxseed diet (Table 2). Serum levels of sex hormone binding globulin were essentially unchanged in normal and overweight/obese women.

Table 2: Effects of Dietary Flaxseed on Serum Sex Hormone Levels among Overweight/Obese Women

	Baseline	Change from Baseline to Follow-		Change from Baseline to Follow-	
	Level		up 1		up 2
Hormone	mean	mean	95% CI	mean	95%CI
Testosterone (pg/ml)	180.2	-13.4 *	-26.5, -0.3	-7.5	-18.2, 3.2
Estrone (pg/ml)	36.0	-7.0 *	-13.4, -0.5	-6.4*	-11.5, -1.3
Estradiol (pg/ml)	12.3	-3.4	-9.0, 2.1	-3.3	-8.9, 2.3

**p*≤..05

These data suggest that dietary flaxseed may have favorable effects on serum estrogen and androgen levels in overweight/obese women. Thus, dietary flaxseed may be beneficial in reducing the risk of breast cancer. Larger studies with a separate control group are needed to confirm these findings. Statistical analyses are currently underway to examine the relationship between dietary flaxseed intake and changes in urinary estrogen metabolites.

One study participant notified the Principal Investigator that she was experiencing a vaginal discharge that began during the study that had continued to worsen during the four months after stopping flaxseed. She was concerned that this adverse health event could be related to flaxseed intake. As part of her own routine primary care, she had been worked up for bacterial, fungal and yeast infections and had a PAP smear and pelvic ultrasound which were within normal limit. After notifying the institutional IRB about this reported occurrence and reviewing the published literature, she was informed by the Principal Investigator that there is no available evidence to suggest that this event would be related to the ingestion of flaxseed.

Key Research Accomplishments:

Preliminary findings indicate that serum levels of estrone, estradiol and testosterone
declined after flaxseed supplementation but these findings were not statistical
significant, and there was no evidence of greater declines with increasing dose and

duration of intake. Additional analyses revealed that dietary flaxseed may have especially favorable effects on serum estrogen and androgen levels in overweight/obese women.

 Urinary specimens were sent to Dr. Mindy Kurzer, and have been analyzed for lignan level (as a measure of compliance) and estrogen metabolites. Statistical analyses are currently underway to examine relation between dietary flaxseed and changes in urinary estrogen metabolites.

Conclusions:

Our preliminary findings suggest that dietary flaxseed may have beneficial effects on serum hormone levels that are important in the etiology of breast cancer, that are more pronounced in overweight/obese women. Additional studies focusing on the direct effects on flaxseed on breast tissue appear warranted. Additional larger studies with a control group would also be desirable.

Contractual Issues

After discussion with the CORE, no additional justification of the 36-month performance period is necessary.

Reportable Outcomes:

An application submitted to the Americian Institute of Cancer Research was funded to examine the effects of dietary flaxseed on other blood biomarkers of cancer risk (i.e., insulin-like growth factor, insulin-like growth factor binding protein, c-peptide) using remaining blood specimens from this study. The title of this proposal is "Intervention Study of Flaxseed in Postmenopausal Women: Effects on Biomarkers of Cancer Risk". After discussion with the COR, the grant number is not necessary as this is not a DOD grant.

References: Not Applicable

Appendices: Investigators and Bibliography/Posters

FINAL REPORT

BIBLIOGRAPHY/POSTERS

Heersink J. Serum testosterone, estrone, estradiol, and sex-hormone binding blogulin responses in healthy postmenopausal women participating in a flaxseed intervention study. Master's thesis presented at the University of Massachusetts at Amherst, 2004

Sturgeon SR, Heersink JL, Bertone-Johnson ER, Stanczyck F, Sabelawski S, Kurzer M, Volpe S, and Bigelow C. Effects of dietary flaxeed on serum levels of estrogens and androgens in postmenopausal women. Poster presented at Era of Hope, Department of Defense Breast Cancer Research Program Meeting. June 8-11, 2005, Philadelphia, Pennsylvania.

Sturgeon SR, Heersink JL, Bertone-Johnson ER, Stanczyck F, Sabelawski S, Kurzer M, Volpe S, and Bigelow C. Effects of dietary flaxeed on serum levels of estrogens and androgens in postmenopausal women. (manuscript in preparation)

Sturgeon SR, Bertone-Johnson ER, Stanczyck F, Sabelawski S, Kurzer M, Volpe S, and Bigelow C. Effects of dietary flaxeed on urinary levels of estrogen metabolites in postmenopausal women. (manuscript in preparation)

LIST OF PERSONNEL

Susan R. Sturgeon, Dr.PH, MPH Carol Bigelow, Ph.D. Stella Volpe, Ph.D. Sara Sabelawski, MS Joanna Heersink, Ph.D. Frank Stanczyck, Ph.D. Mindy Kurzer, Ph.D.